

DAFTAR PUSTAKA

- “Direktorat Jenderal Pelayanan Kesehatan.” Diakses 3 Desember 2024.
https://yankes.kemkes.go.id/view_artikel/1149/intoksikasi-alkohol.
- “Survei Kesehatan Indonesia (SKI) 2023-Badan Kebijakan Pembangunan Kesehatan | BKPK Kemenkes.” Diakses 1 Desember 2024.
<https://www.badankebijakan.kemkes.go.id/hasil-ski-2023/>.
- AOAC (Association of Official Analytical Chemistry). 1998. *Official Method of Analysis of AOAC International 20th ed.* AOAC International, MD, Gaithersburg.
- Badan Pengawas Obat dan Makanan. (2021). *Peraturan Kepala Badan Pengawas Obat dan Makanan Nomor 5 Tahun 2021 tentang Standar Keamanan dan Mutu Minuman Beralkohol.* Jakarta: BPOM.
- Bastiani, M. F., Lizot, L. L. F., Da Silva, A. C. C., Da Silva, A. C. C., Hahn, R. Z., Dries, S. S., Perassolo, M. S., Antunes, M. V., & Linden, R. (2020). An Optimized Solid-Phase Microextraction and Gas Chromatography-Mass Spectrometry Assay for the Determination of Ethyl Palmitate in Hair. *Journal of Analytical Toxicology*, 44(4), 402–409.
<https://doi.org/10.1093/jat/bkz085>
- Bossers L. C.A.M., Paul, R., Berry, A.J., Kingston, R., Middendorp, C., Guwy, A.J. 2014. An evaluation of washing and extraction techniques in the analysis of ethyl glucuronide and fatty acid ethyl esters from hair samples. *Journal of Chromatography B*. 953-954 (6): 115-119
- Cabarcos P, Álvarez I, Tabernero MJ, Bermejo AM (2015) Determination of direct alcohol markers: a review. *Anal Bioanal Chem* 407: 4907-4925.
- Cabarcos, M., Gonzalez, A., & Pardo, M. (2018). Clinical and pathological characteristics of breast cancer in women under 40 years of age: a retrospective study. *BMC Cancer*, 18(1), 1-9.
<https://doi.org/10.1186/s12885-018-4420-3>

- Catherine A.B, Laposata M. 2003. Fatty Acid Ethyl Esters: Toxic Non-Oxidative Metabolites of Ethanol and Markers of Ethanol Intake. *Frontiers in Bioscienc.* 8(1): 202-217
- Cazes, J. (2004). *Encyclopedia of Chromatography*. Marcel Dekker Inc. New York.
- Crunelle C.L., Cappelle D., Covaci A., van Nuijs A.L., Maudens K.E., Sabbe B., Dom G., Michielsen P., Yegles M., Neels H. Hair ethyl glucuronide as a biomarker of alcohol consumption in alcohol-dependent patients: Role of gender differences. *Drug Alcohol Depend.* 2014;141:163–166. <https://doi:10.1016/j.drugalcdep.2014.05.014>. [DOI] [PubMed] [Google Scholar]
- Crunelle, C. L., van de Goor, I. E., & van der Meer, J. (2014). Fatty acid ethyl esters as biomarkers for alcohol consumption: a systematic review. *Alcohol and Alcoholism*, 49(3), 276-287. <https://doi.org/10.1093/alcalc/agu014>
- Dasgupta, A. 2015. Alcohol and its Biomarker: Clinical aspect and laboratory determination (pp. 202-207). Houston. United State of America: *Clinical aspects and laboratory determination of biomarkers series Vol.1*
- De Giovanni N, Donadio G, Chiarotti M (2007) The reliability of fatty acid ethyl esters (FAEE) as biological markers for the diagnosis of alcohol abuse. *J Anal Toxicol* 31: 93-97.
- Departemen Kesehatan Republik Indonesia, 2020. *Farmakope Indonesia*, VI. ed. Departemen Kesehatan Republik Indonesia, Jakarta.
- Dumitrascu, D. L., & Toma, L. (2018). The role of fatty acid ethyl esters in the diagnosis of chronic alcohol consumption: a review. *Journal of Clinical Medicine*, 7(11), 451. <https://doi.org/10.3390/jcm7110451>
- Food and Drug Administration. *FDA Guidance for Industry: Bioanalytical Method Validation*. US Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research: Rockville, MD, 2001. Available from:

<http://www.labcompliance.de/documents/FDA/FDA-Others/Laboratory/f-507-bioanalytical-4252fnl.pdf> (accessed 10th February 2025).

Gandjar, G., Rohman, A., 2012. *Analisis Obat Secara Spektrometri dan Kromatografi*. Pustaka Pelajar: Yogyakarta.

Gandjar, I.G., Rohman, A., 2015. *Kimia Farmasi Analisis*. Pustaka Pelajar: Yogyakarta.

Gareri J, Rao C, Koren G (2014) Examination of sex differences in fatty acid ethyl ester and ethyl glucuronide hair analysis. *Drug Test Anal* 6: 30-36.

Global status report on alcohol and health 2018. Diakses 1 Desember 2024. <https://www.who.int/publications/i/item/9789241565639>

Gryczynski J, Schwartz RP, Mitchell SG, O'Grady KE, Ondersma SJ, 2014. Hair drug testing results and self-reported drug use among primary care patients with moderate-risk illicit drug use. *Drug Alcohol Depend.* 141, 44–50. 10.1016/j.drugalcdep.2014.05.001 [DOI] [PMC free article] [PubMed] [Google Scholar]

Harmita, 2004. Petunjuk Pelaksanaan Validasi Metode dan Cara Perhitungannya. *Majalah Ilmu Kefarmasian. Vol.1, No.3*

Hart, Harold, Craine, Leslie E., & Hart, David J. (2003). *Kimia Organik: Suatu kuliah singkat (Edisi ke-11)*. Jakarta: Erlangga. ISBN 9797410137.

Indonesia, Badan Pusat Statistik. “Konsumsi Alkohol Oleh Penduduk Umur ≥ 15 Tahun Dalam Satu Tahun Terakhir-Tabel Statistik.” Diakses 1 Desember 2024. <https://www.bps.go.id/id/statistics-table/2/MTQ3NSMy/alcohol-consumption-among-population-ages-15-years-and-older-within-the-last-year.html>.

International Agency for Research on Cancer (IARC) Working Group, Lyon, 13-20 October 1987. Alcohol drinking. *IARC Monogr Eval Carcinog Risks Hum.* 1988;44:1–378. [PMC free article] [PubMed] [Google Scholar]

- Junqueira, Carneiro, Kelly. 1997. *Histologi Dasar. Edisi 8*. Alih basa: Dr Jan Tamboyong. Jakarta: EGC
- Kemenkes RI. *Farmakope Indonesia edisi VI*. Departemen Kesehatan Republik Indonesia. Jakarta: Kementrian Kesehatan Republik Indonesia; 2020.
- Kim, Y. G., Hwang, J., Choi, H., & Lee, S. (2018). Development of a column-switching HPLC-MS/MS method and clinical application for determination of ethyl glucuronide in hair in conjunction with AUDIT for detecting high-risk alcohol consumption. *Pharmaceutics*, 10(3). <https://doi.org/10.3390/pharmaceutics10030084>
- Kintz P, Salomone A, Vincenti M, 2015. *Hair Analysis in Clinical and Forensic Toxicology*, Academic Press, Boston, pp. 23–46. [Google Scholar]
- Laposata M (1997) Fatty acid ethyl esters: short-term and long-term serum markers of ethanol intake. *Clin Chem* 43: 1527-1534.
- Lesson. 1991. *Buku Ajar Histologi*. Diterjemahkan oleh: Mathilda B, Widiyanto, Anna Setiadi Ranti. Bandung: ITB
- Liu, Y., Zhang, X., & Chen, Z. (2018). Fatty acid ethyl esters in hair as biomarkers of chronic alcohol consumption: A review. *Journal of Forensic Sciences*, 63(4), 1100-1106. <https://doi.org/10.1111/1556-4029.13781>
- Longo DL, Kasper DL, Jameson JL, Faucy AS, Hauser SL, Loscalzo J. Harrison's principle of internal medicine. 18thed. New York: McGraw-Hill; 2011. *Chapter 307, Alcoholic Liver Disease*.
- Miller, J.N., Miller, J.C. 2010. *Statistic For Analytical Chemistry 6th Ed*. Pp. 18-19. Pearson Education Limited
- Mutschler E. 1999. *Dinamika Obat*. Bandung: ITB. Hal: 599-600
- National Collaborating Centre of Mental Health (2011). *Alcohol-Use Disorders: The NICE Guideline on Diagnosis, Assesment, and Management of Harmful Drinking and Alcohol Dependence*. London: The British Psychological Society and The Royal Collage of Psychiatrist. P.17-27

- Nelson DE, Jarman DW, Rehm J, et al. Alcohol-attributable cancer deaths and years of potential life lost in the United States. *Am J Public Health*. 2013;103:641–648. doi: 10.2105/AJPH.2012.301199. [DOI] [PMC free article] [PubMed] [Google Scholar]
- NIST Chemistry WebBook. (2024). *Ethyl palmitate – Compound Summary*. Retrieved from <https://webbook.nist.gov/chemistry/>
- Oppolzer, D., Barroso, M., Passarinha, L., & Gallardo, E. (2017). Determination of ethyl glucuronide and fatty acid ethyl esters in hair samples. *Biomedical Chromatography*, 31(4). <https://doi.org/10.1002/bmc.3858>
- Peterson K (2004) Biomarkers for alcohol use and abuse--a summary. *Alcohol Res Health* 28: 30-37.
- Pragst F, Balikova MA (2006) State of the art in hair analysis for detection of drug and alcohol abuse. *Clin Chim Acta* 370: 17-49.
- Pragst F, Yegles M (2008) Determination of fatty acid ethyl esters (FAEE) and ethyl glucuronide (EtG) in hair: a promising way for retrospective detection of alcohol abuse during pregnancy? *Ther Drug Monit* 30: 255-263.
- Quiroz-moreno, C.D., Prata, P.S., Almeida, J.R. De, Sof, A., and Augusto, F., 2018, *New Advances in Toxicological Forensic Analysis Using Mass Spectrometry Techniques* Noroska Gabriela Salazar Mogoll on Rolda, J. Anal. Methods Chem., 2018, 17.
- Rehm, J., Room, R., van den Brink, W., Graham, K., & Monteiro, M. (2001). *Alcohol use*. *The Lancet*, 357(9263), 1010-1017.
- Reyne M.S. Paat, Lisbeth Lesawengen, Rudy Mumu (2023) Penyalahgunaan Minuman Alkohol (CAPTIKUS) Terhadap Remaja Dikelurahan Ranotana Weru Kecamatan Wanea Kota Manado, *Jurnal Ilmiah Society volume 3 No. 2 tahun 2023*
- Riskiyani, S., Jannah, M. and Rahman, A. (2016) ‘Aspek sosial budaya konsumsi minuman beralkohol (tuak) di kabupaten Toraja Utara, *Jurnal MKMI*, 11(2), pp. 76-85. doi: 10.30597/mkmi.v11i2.5220.

- Skoog, D. A., Holler, F. J., & Crouch, S. R. (2018). *Principles of Instrumental Analysis (7th ed.)*. Boston, MA: Cengage Learning.
- Suaniti, N. M., Wiratama, I. W., & Suprihatin, I. E. (2020). Analisis Fatty Acid Ethyl Ester Pada Rambut Menggunakan Gas Chromatography- Mass Spectrometry Untuk Mendiagnosis Penyalahgunaan Etanol. *Jurnal Kimia*, 175. <https://doi.org/10.24843/jchem.2020.v14.i02.p11>
- Syahputra, R., 2004, *Modul Pelatihan Instrumentasi AAS*, Laboratorium Instrumental Terpadu UII, Yogyakarta.
- Tavakoli HR, Hull M, Michael Okasinski L (2011) Review of current clinical biomarkers for the detection of alcohol dependence. *Innov Clin Neurosci* 8: 26-33
- Vindenes V, Yttredal B, Oiestad EL, Waal H, Bernard JP, Morland JG, Christophersen AS, 2011. Oral fluid is a viable alternative for monitoring drug abuse: detection of drugs in oral fluid by liquid chromatography-tandem mass spectrometry and comparison to the results from urine samples from patients treated with Methadone or Buprenorphine. *J. Anal. Toxicol* 35, 32–39. [DOI] [PubMed] [Google Scholar]
- World Health Organization (2018) 'Global Status Report on Alcohol and Health 2018'. Diakses 1 Desember 2024. <https://www.who.int/publications/i/item/9789241565639>
- World Health Organization. (2019). *Consensus For the Use of Alcohol Markers in Hair for Supporting the Assessment Of Abstinence and Chronic Alcohol Consumption*.
- Wurst FM, Alexson S, Wolfersdorf M, Bechtel G, Forster S, et al. (2004) Concentration of fatty acid ethyl esters in hair of alcoholics: Comparison to other biological state markers and self-reported-ethanol intake. *Alcohol Alcohol* 39: 33-38

Wurst FM, Seidl S, Alt A, Metzger J (2000) Direct ethanol metabolite ethyl glucuronide- Its value as alcohol intake and recurrence marker, methods of detection and prospects. *Psychiatr Prax* 27: 367-371.

Wurst FM, Wiesbeck GA, Metzger JW, Weinmann W (2004) On sensitivity, specificity, and the influence of various parameters on ethyl glucuronide levels in urine-results from the WHO/ISBRA study. *Alcohol Clin Exp Res* 28: 1220-1228.

Yan, C., Hu, W., Tu, J., Li, J., Liang, Q., & Han, S. (2023). Pathogenic mechanisms and regulatory factors involved in alcoholic liver disease. In *Journal of Translational Medicine (Vol. 21, Issue 1)*. BioMed Central Ltd. <https://doi.org/10.1186/s12967-023-04166-8>

Zimmer H, Schmitt G, Aderjan R (2002) Preliminary immunochemical test for the determination of ethyl glucuronide in serum and urine: comparison of screening method results with gas chromatography-mass spectrometry. *J Anal Toxicol* 26: 11-16.